

Instanton–anti–instanton molecule with non–zero–modes of quarks included

A.G.Zubkov, O.V.Dubasov, B.O.Kerbikov,
Institute of Theoretical and Experimental Physics
117218, Moscow, B.Chermushkinskaya 25, Russia
E-mail:borisk@heron.itep.ru

Abstract

A straightforward algebraic derivation is carried out of light quark propagator and effective action in the instanton–anti–instanton molecule. Exact expressions are obtained which contain contributions of all quark modes. Possible implications of the results include chiral phase transition and correlation functions of mesons and baryons.

1 Introduction

The theory of light quarks in the instanton vacuum has a long history – see e.g. review papers [1-3]. Recently a new impetus to the instanton liquid model¹ has been provided by lattice measurements of quark and gluon correlators in coordinate space [4] showing the important role of instantons in forming hadronic wave functions and mass spectra. The standard theory of the instanton vacuum is based on the assumption of quark zero–modes dominance. This assumption was questioned in [5], and it was argued that non-zero–modes of each instanton play an important role in addition to zero–modes. To our knowledge it was the first systematic treatment of higher modes although their role in specific effects and within various approximations were considered in some earlier papers [6-8].

Here we wish to present a complete treatment with all modes included of the so–called instanton–anti–instanton molecule, i.e. the $I\bar{I}$ pair ”bound”

¹Sometimes also called instanton gas model

through light quarks. This object plays an important role in the instanton model of the QCD vacuum. According to the standard scenario the chiral symmetry restoration occurs due to the onset of the molecular component in the instanton–anti–instanton ensemble [9-15]. The analysis of chiral phase transition is not the purpose of the present paper. Our main objective is the detailed calculation of the molecule including quark propagator and effective action. Toolled with these results one can reexamine the chiral symmetry restoration in more details. Another even more immediate implication is the study of interactions in mesons and baryons induced by correlated $I\bar{I}$ pairs [14]. The corresponding effective Lagrangian is derived from the quark propagator. As we shall see the quark propagator in the quantity in which the effects of non–zero–modes are mostly pronounced.

The systematic study of $I\bar{I}$ molecule was performed in 1979 by C.Lee and W.A.Bardeen [16]. The propagator and the effective action were constructed making use of the power series in two independent parameters $m\rho$ and ρ/d , where m is the quark dynamical mass, ρ is the size of (anti–) instanton, d –the distance between the two centers. In the propagator the interference terms between zero–modes and continuum were taken into account in the leading order with respect to the above parameters. Using this propagator the quark effective action was constructed. Surprisingly enough it contains only zero–modes overlap. In other words it was shown that the propagator is more affected by non-zero–modes than the effective action. The overlap of zero–modes in the effective action describes logarithmic attraction between instanton and anti–instanton.

The revival of the interest to the role of non-zero–modes on one hand and to the $I\bar{I}$ molecule on the other calls for the reexamination of the problem. Below we develop an approach different from that of [16]. In its spirit our formalism is close to the multiple scattering theory [17,13], or scattering theory in a system with confinement [18]. Along these lines we have found the exact solution of the problem with all modes included. The final answers for the quark propagator and effective action are given in terms of matrix elements involving individual (anti–) instanton zero–modes and individual propagators with zero–modes deleted. Unfortunately the corresponding integrals may not be evaluated analytically and we estimate them by the orders of magnitudes with respect to the parameters ρ/d and $m\rho$. In this way we present the non–zero–modes corrections to the effective action evaluated by Lee and Bardeen [16]. To get accurate results one has to perform numerical calculations which is beyond the scope of this paper.

The outline of this work is as follows. In Section 2 we introduce notations and derive the "two-centers" equation for the quark propagator in $I\bar{I}$ molecule. Section 3 is devoted to the evaluation of the quark propagator in $I\bar{I}$ molecule. We present an exact expression for the propagator and an approximate form in which higher modes contribution is accounted for in the leading order. We argue that the interference of zero modes with continuum is generally of the same order as zero modes contribution. Section 4 deals with fermionic effective action. Again we find an exact answer with the contributions of all modes included. In Section 5 we discuss estimates for the propagator and for the effective action based on the existence of two small parameters – the ratio of the instanton size to the size of the molecule and the quark current mass. Finally in Section 6 we discuss our results and suggest various ways of their implementations.

2 Basic definitions and the general approach

As usual in the instanton physics we work in Euclidean metric. Also in line with the common lore we use the superposition ansatz for the $I\bar{I}$ system

$$A_\mu(x) = \sum_{i=1,2} A_\mu^{(i)}(x - R_i) = A_\mu^I(x) + A_\mu^{\bar{I}}(x - d), \quad (1)$$

where $i = 1$ corresponds to I placed at the origin ($R_1 \equiv R_I = 0$), $i = 2$ corresponds to $\bar{I}$ placed at $R_2 \equiv R_{\bar{I}} = d$. In singular gauge we can write

$$gA_\mu^I(x) = \frac{\bar{\eta}_{a\mu\nu} x_\nu \rho^2 \Omega_I^+ \tau_a \Omega_I}{x^2(x^2 + \rho^2)}, \quad (2)$$

where ρ and Ω_I are the size and color orientation of the instanton. Anti-instanton is assumed to have the same size ρ (this is not crucial), and its field follows from (2) though the substitution $\bar{\eta} \rightarrow \eta$ and $x \rightarrow (x - d)$. Next we introduce the free propagator S_0 , the single (anti-) instanton propagator S_i with $i = 1$ for I and $i = 2$ for $\bar{I}$, and the total propagator S in the field (1):

$$\begin{aligned} S_0^{-1} &= (-i\hat{\partial} - im), \\ S_i^{-1} &= (-i\hat{\partial} - g\hat{A}^{(i)} - im), \\ S^{-1} &= (-i\hat{\partial} - g\hat{A} - im). \end{aligned} \quad (3)$$

Note that our free propagator S_0 includes the current quark mass m which is not the case in Lee and Bardeen paper [16].

The instanton propagator has a formal spectral representation over normal modes

$$S_I = \sum_{n=0}^{\infty} \frac{|u_n^I \rangle \langle u_n^I|}{\lambda_n - im}, \quad (4)$$

with eigenfunctions $|u_n^I \rangle$ satisfying the equation

$$(-i\hat{\partial} - g\hat{A}^I)|u_n^I \rangle = \lambda_n |u_n^I \rangle. \quad (5)$$

Of special role are zero-modes with $\lambda_0 = 0$ which have negative chirality

$$\gamma_5 |u^I \rangle = -|u^I \rangle, \quad (6)$$

where index $n = 0$ is omitted. Anti-instanton zero-modes $|u^{\bar{I}} \rangle$ have positive chirality. Zero-mode has the following explicit form

$$\begin{aligned} (u^I(x))_{ci} &= \frac{\rho}{\pi(2x^2)^{1/2}(x^2 + \rho^2)^{3/2}} \left[\left(\frac{1 - \gamma_5}{2} \right) \hat{x} \right]_{ij} \varepsilon_{jc'} \Omega_{c'}^+ = \\ &= \frac{\rho}{\pi(2x^2)^{1/2}(x^2 + \rho^2)^{3/2}} (\tau_x)_{ij} \varepsilon_{jc'} \Omega_{c'}^+, \end{aligned} \quad (7)$$

where c and i are color and Lorentz indices, ε is antisymmetric tensor, $\tau_x = \tau_\mu x_\mu$, $\tau_\mu = (\vec{\tau}, i)$, $\tau_\mu^+ = (\vec{\tau}, -i)$ with $\vec{\tau}$ being the Pauli matrix. For anti-instanton x is replaced by $(x - d)$, $(1 - \gamma_5) \rightarrow (1 + \gamma_5)$, $\tau_x \rightarrow \tau_{x-d}^+$.

The non-zero-modes part of the instanton propagator (3-4) is known explicitly in the chiral limit $m \rightarrow 0$ [19] and given by [19,6,7]

$$\begin{aligned} S'_I(x, y) &= \sum_{n>0} \frac{u_n^I(x)(u_n^I(y))^+}{\lambda_n} = \\ &= G_0(x - y) \left(1 + \frac{\rho^2}{x^2 y^2} \tau_x \tau_y^+ \right) (\pi_x \pi_y)^{-1/2} - \\ &\quad - \frac{\rho^2 (\pi_x \pi_y)^{-1/2}}{4\pi^2 x^2 y^2 (x - y)^2} \tau_x \left[\frac{\hat{\tau}^+ \tau_{x-y}}{x^2 + \rho^2} \left(\frac{1 + \gamma_5}{2} \right) + \frac{\tau_{x-y}^+ \hat{\tau}}{y^2 + \rho^2} \left(\frac{1 - \gamma_5}{2} \right) \right] \tau_y, \end{aligned} \quad (8)$$

where $G_0(x - y)$ is the free propagator of massless fermion (not to be confused with S_0 given by (3) in which the quark current mass is included)

$$G_0(z) = -\frac{i}{2\pi^2} \frac{\hat{z}}{z^4}, \quad (9)$$

and

$$\pi_z = 1 + \rho^2/z^2. \quad (10)$$

The propagator $S'_I(x, y)$ obeys the Green's function equation with $m = 0$ and zero-mode deleted

$$(-i\hat{\partial} - g\hat{A}^I)S'_I = \mathbf{1} - |u^I \rangle \langle u^I|. \quad (11)$$

Mass corrections of the order $0(m)$ to S'_I can be easily found from the spectral representation (4) rewritten in the form

$$S_I = \frac{|u^I \rangle \langle u^I|}{-im} + \sum_{n>0} \frac{\lambda_n}{\lambda_n^2 + m^2} |u_n^I \rangle \langle u_n^I| + im \sum_{n>0} \frac{|u_n^I \rangle \langle u_n^I|}{\lambda_n^2 + m^2}.$$

In the chiral limit the second term in the right-hand side yields S'_I (see the first line in (8)). To get $0(m)$ correction from the third term we note that due to the orthogonality of eigenmodes

$$\sum_{n>0} \frac{|u_n^I \rangle \langle u_n^I|}{\lambda_n^2} = \left(\sum_{n>0} \frac{|u_n^I \rangle \langle u_n^I|}{\lambda_n} \right)^2 = (S'_I)^2.$$

Thus the quark propagator in the single instanton field up to order m^2 terms has form [6]

$$\begin{aligned} S_I(x, y) &= \frac{u^I(x)(u^I(y))^+}{-im} + S'_I(x, y) + \\ &+ im \int dz S'_I(x, z) S'_I(z, y). \end{aligned} \quad (12)$$

We note in passing that the instanton liquid model (see [2,3] and references therein) relies on the following approximation for the quark propagator in the single instanton field

$$S_I = \frac{|u^I \rangle \langle u^I|}{-im} + (-i\hat{\partial})^{-1}. \quad (13)$$

The validity of this ansatz in instanton liquid model was discussed in [5]. The situation for the $I\bar{I}$ molecule will be elucidated in what follows.

For further purposes it will be convenient to recast (12) into the following form

$$S_I = \frac{|u^I \rangle \langle u^I|}{-im} + S_0 + \Delta_I, \quad (14)$$

so that

$$\begin{aligned}\Delta_I \equiv \Delta_1 &= S'_I - S_0 + im(S'_I)^2 = (S'_I - G_0) + im[(S'_I)^2 - G_0^2] \equiv \\ &\equiv \delta_1 + im[(S'_I)^2 - G_0^2],\end{aligned}\quad (15)$$

where G_0 is given by (9). For $\bar{I}$ similar notations are introduced with $\Delta_{\bar{I}} \equiv \Delta_2$.

Now we derive the two-centers equation for the propagator S defined by (3). It follows straightforwardly from the superposition ansatz (1) and the definitions (3). Namely one can write

$$\begin{aligned}S &= \frac{1}{S_1^{-1} + S_2^{-1} - S_0^{-1}} = S_2 \frac{1}{S_1(S_1^{-1} + S_2^{-1} - S_0^{-1})S_2} S_1 = \\ &= S_2 \frac{1}{S_0 - (S_1 - S_0)S_0^{-1}(S_2 - S_0)} S_1 = S_1 \frac{1}{S_0 - (S_2 - S_0)S_0^{-1}(S_1 - S_0)} S_2.\end{aligned}\quad (16)$$

Generalization for more than two centers may be found in [13]. Next task is the actual calculation of (16), i.e. the inversion of the operator in the denominator. This will be done exactly without resorting to the approximations of the type

$$S \simeq S_0 + \sum_i (S_i - S_0).$$

3 Quark propagation function in the molecule

Using (14) the propagator (16) may be rewritten as

$$S = S_2 Z^{-1} S_1, \quad (17)$$

$$\begin{aligned}Z &= S_0 - \Delta_1 S_0^{-1} \Delta_2 - \frac{|u^1 \rangle \langle u^1|}{-im} S_0^{-1} \Delta_2 - \Delta_1 S_0^{-1} \frac{|u^2 \rangle \langle u^2|}{-im} + \\ &+ |u^1 \rangle \frac{V_{12}}{m^2} \langle u^2|,\end{aligned}\quad (18)$$

$$V_{12} = \langle u^1 | S_0^{-1} | u^2 \rangle = \langle u^1 | -i\hat{\partial} | u^2 \rangle = \langle u^1 | g\hat{A}^{(2)} | u^2 \rangle. \quad (19)$$

Performing calculations one may use any of the two equivalent expressions (16) for S . At intermediate stages of calculations the $1 \leftrightarrow 2$ permutation symmetry is lost as in (18). In final answers this symmetry is restored.

To find Z^{-1} we will use the technique based on the relation

$$\frac{1}{A + |\zeta > a < \eta|} = \frac{1}{A} - \frac{1}{A} |\zeta > \frac{a}{1 + \langle \eta | \frac{a}{A} | \zeta >} < \eta | \frac{1}{A}, \quad (20)$$

with A being an operator and a – c -number. Applying it to (18) one gets

$$\begin{aligned} Z^{-1} &= \frac{1}{W + |u^1 > \frac{V_{12}}{m^2} < u^2|} = \\ &= \frac{1}{W} - \frac{1}{W} |u^1 > \frac{V_{12}}{m^2 + V_{12} < u^2 | W^{-1} | u^1 >} < u^2 | \frac{1}{W}, \end{aligned} \quad (21)$$

$$W = K - \frac{|u^1 > < u^1|}{-im} S_0^{-1} \Delta_2 - \Delta_1 S_0^{-1} \frac{|u^2 > < u^2|}{-im}, \quad (22)$$

$$K = S_0 - \Delta_1 S_0^{-1} \Delta_2. \quad (23)$$

To find W^{-1} one has to apply equation (20) repeatedly twice. After some straightforward but tedious work of collecting all terms in (21) we arrive at

$$\begin{aligned} Z^{-1} &= K^{-1} + \frac{im + \alpha}{D} K^{-1} |u^1 > < u^1 | R_2 K^{-1} + \frac{im + \beta}{D} K^{-1} R_1 |u^2 > < u^2 | K^{-1} - \\ &\quad \frac{V_{12} + A_{12}}{D} K^{-1} |u^1 > < u^2 | K^{-1} - \frac{V_{21} + A_{21}}{D} K^{-1} R_1 |u^2 > < u^1 | R_2 K^{-1}, \end{aligned} \quad (24)$$

where

$$\begin{aligned} \alpha &= < u^2 | K^{-1} \Delta_1 S_0^{-1} | u^2 >, \\ \beta &= < u^1 | S_0^{-1} \Delta_2 K^{-1} | u^1 >, \\ V_{12} &= < u^1 | S_0^{-1} | u^2 >, V_{21} = < u^2 | S_0^{-1} | u^1 >, \\ A_{12} &= < u^1 | S_0^{-1} \Delta_2 K^{-1} \Delta_1 S_0^{-1} | u^2 >, \\ A_{21} &= < u^2 | S_0^{-1} \Delta_1 \tilde{K}^{-1} \Delta_2 S_0^{-1} | u^1 >, \\ K &= S_0 - \Delta_1 S_0^{-1} \Delta_2, \quad \tilde{K} = S_0 - \Delta_2 S_0^{-1} \Delta_1, \\ D &= m^2 + (V_{12} + A_{12})(V_{21} + A_{21}) - im(\alpha + \beta) - \alpha\beta. \\ R_i |u^j > &= \Delta_i S_0^{-1} |u^j >, \quad < u^j | R_i = < u^j | S_0^{-1} \Delta_i, \quad i, j = 1, 2. \end{aligned} \quad (25)$$

The final step is to insert (24) into (17) and use (14) for S_i . This yields the following expression for the propagator

$$S = \frac{1}{2} (S_0 + \Delta_2) K^{-1} (S_0 + \Delta_1) + \frac{1}{2} (S_0 + \Delta_1) \tilde{K}^{-1} (S_0 + \Delta_2) +$$

$$\begin{aligned}
& + \frac{im + \alpha}{D} (S_0 + \Delta_2) K^{-1} |u^1 \rangle \langle u^1| \tilde{K}^{-1} (S_0 + \Delta_2) + \\
& + \frac{im + \beta}{D} (S_0 + \Delta_1) \tilde{K}^{-1} |u^2 \rangle \langle u^2| K^{-1} (S_0 + \Delta_1) - \\
& - \frac{V_{12} + A_{12}}{D} (S_0 + \Delta_2) K^{-1} |u^1 \rangle \langle u^2| K^{-1} (S_0 + \Delta_1) - \\
& - \frac{V_{21} + A_{21}}{D} (S_0 + \Delta_1) \tilde{K}^{-1} |u^2 \rangle \langle u^1| \tilde{K}^{-1} (S_0 + \Delta_2). \tag{26}
\end{aligned}$$

This expression is an exact one. It relies only on the standard superposition assumption (1) for the molecular field and on the formal representation (14) for the individual instanton propagator. All individual (anti-) instanton modes are included. No any small parameter like ρ/d or $m\rho$ was used in the derivation. To get from (26) an insight into the physical picture one should either perform numerical calculations or resort to approximations and estimates. This last strategy will be used in what follows.

Keeping in (26) only linear terms in Δ_i one obtains

$$\begin{aligned}
S \simeq S_0 + \Delta_1 + \Delta_2 + \frac{im + \alpha}{d} (1 + R_2) |u^1 \rangle \langle u^1| (1 + R_2) + \\
+ \frac{im + \beta}{d} (1 + R_1) |u^2 \rangle \langle u^2| (1 + R_1) - \\
- \frac{V_{12}}{d} (1 + R_2) |u^1 \rangle \langle u^2| (1 + R_1) - \frac{V_{21}}{d} (1 + R_1) |u^2 \rangle \langle u^1| (1 + R_2), \tag{27}
\end{aligned}$$

where

$$d = m^2 + V_{12}V_{21} - im(\alpha + \beta). \tag{28}$$

This approximation coincides with the result of Lee and Bardeen [16]. In the chiral limit, more precisely when $m^2 \ll V_{12}V_{21}$, one gets even simpler expression

$$\begin{aligned}
S \simeq S_0 + \Delta_1 + \Delta_2 - V_{21}^{-1} (1 + R_2) |u^1 \rangle \langle u^2| (1 + R_1) - \\
- V_{12}^{-1} (1 + R_1) |u^2 \rangle \langle u^1| (1 + R_2). \tag{29}
\end{aligned}$$

It is tempting to simplify (29) further and neglect Δ_i and R_i (see e.g. [3,14]). However such approximation is not justified since as it was already mentioned in [16] and will be discussed below in Section 5 all terms in (29) are in general of the same order of magnitude.

4 Fermionic effective action

We define the quark effective action in $I\bar{I}$ molecule as

$$\exp\{\Gamma_{eff}\} = \frac{\det S^{-1}}{\det S_0^{-1}} \quad (30)$$

We leave aside at least temporarily the questions of normalization and regularization – see, e.g. [20-22]. From (16) we get

$$\det S^{-1} = \det S_1^{-1} \cdot \det S_2^{-1} \cdot \det Z \quad (31)$$

where the quantity Z is given by Eq. (18) of the previous section. To calculate $\det Z$ use can be made of an equation similar to (20)

$$\det(A + |\zeta\rangle\langle a|) = \det A \cdot (1 + \langle a|\frac{a}{A}|\zeta\rangle). \quad (32)$$

It is a simple but tedious task to apply (31) repeatedly to (18). In fact no new calculations has be done since the needed quantity essentially coincides with D defined by (24-25). Precisely

$$\det Z = \det S_0 \cdot \det(1 - S_0^{-1}\Delta_1 S_0^{-1}\Delta_2) \frac{D}{m^2}. \quad (33)$$

Combining (30), (31) and (33) one obtains

$$\exp\{\Gamma_{eff}\} = (\frac{\det S_1^{-1}}{\det S_0^{-1}})(\frac{\det S_2^{-1}}{\det S_0^{-1}})\det(1 - S_0^{-1}\Delta_1 S_0^{-1}\Delta_2)D \quad (34)$$

If we now perform standard regularization and normalization procedures [20-22] the final result for the effective action would be

$$\begin{aligned} \Gamma_{eff} = & \Gamma_{eff}^I + \Gamma_{eff}^{\bar{I}} + \ln \det_{reg}(1 - S_0^{-1}\Delta_1 S_0^{-1}\Delta_2^{-1}) + \\ & + N_f \ln[m^2 + (V_{12} + A_{12})(V_{21} + A_{21}) - im(\alpha + \beta) - \alpha\beta]. \end{aligned} \quad (35)$$

This result is an exact one and does not rely upon any approximation except for the superposition ansatz (1). The first two terms describe effective actions of individual (anti-) instantons. The third term corresponds to non-zero modes contribution. Approximated by $\simeq -tr(S_0^{-1}\Delta_1 S_0^{-1}\Delta_2)$ it exhibits exponential damping as compared to the last one. This last term is of the main interest. It consists of the individual zero modes contribution and the interference between zero modes and continuum. The well known approximation to this term is $N_f \ln(m^2 + V_{12}V_{21})$ [16]. In the next section we shall estimate corrections to this expression using the exact result (35).

5 Approximations and estimates

In Section 3 the fermion propagator (26) has been determined and in the previous section the fermion induced effective action (35) has been found. However both expressions are not easy to handle due to high-dimensional integrals which cannot be evaluated analytically. Therefore in the present section we shall resort to estimates based on the existence of two independent parameters supposed to be small. These are ρ/d and $m\rho$, where ρ is the (anti-) instanton size, d - the $I\bar{I}$ separation, m - the current mass of the quark. The widely accepted values of ρ and d are: $\rho \simeq 0.3fm, d \simeq 1fm$ [1-4]. The key quantities entering into most integrals (matrix elements) are Δ_i given by (14-15). According to (15) Δ_i is splitted into mass independent part $\delta_i(x, y) = S'_i(x, y) - G_0(x, y)$ and a contributions proportional to m . Therefore corrections with respect to ρ/d and $m\rho$ are independent.

Let us start by providing arguments in favor of the statement made after equation (29) for the propagator, namely that zero modes interference with continuum is of the same order as zero modes product.

Consider the situation when both x and y are close to the instanton, $x \sim y \sim \rho$. Then using (7) for zero modes one arrives at the following estimate for zero modes product

$$\langle x|u^2 \rangle \langle u^1|y \rangle \sim \rho^{-1}d^{-3}. \quad (36)$$

In the same region the interference term $R_1|u^2 \rangle \langle u^1|$ may be estimated using (8) and (9). This yields

$$\begin{aligned} \langle x|R_1|u^2 \rangle \langle u^1|y \rangle &\simeq \left(\int dz \delta_1(x, z) \partial_z u^2(z - d) \right) \langle u^1|y \rangle \sim \\ &\sim (\rho^4 d^{-3} \rho^{-3}) \rho^{-2} \sim \rho^{-1}d^{-3}. \end{aligned} \quad (37)$$

In (37) we neglected mass corrections and substituted Δ_1 by δ_1 and S_0^{-1} by ∂_z , then performed integration over the shell of size ρ around $z \sim d$ which resulted in ρ^4 volume. Also unjustified (at least in certain regions) is the neglect of Δ_i as compared to S_0 in (29). For example when $x \sim y \sim \rho$ and $(x - y) \sim \rho$ both S_0 and Δ_1 are of the order ρ^{-3} . On the other hand when $x \sim \rho$ and $y \sim d$ zero modes product dominates over interference.

From the above example it is clear that to make estimates one should consider different regions of integration, typically the "near" region $x \sim y \sim \rho$, the "far" region $x \sim y \sim d$, and the "mixed" region $x \sim \rho, y \sim d$.

Looking for the estimate with respect to ρ/d one substitutes S_0^{-1} either by $1/d$, or by $1/\rho$. Attention should be paid to chirality arguments, namely the matrix elements $\langle u^1 | \gamma_\mu \gamma_\nu \dots \gamma_\rho | u^2 \rangle$ and $\langle u^1 | \gamma_\mu \gamma_\nu \dots \gamma_\rho | u^1 \rangle$ are respectively nonzero for odd and even number of γ matrices.

Consider now the essential part of the effective action (35)

$$\tilde{\Gamma}_{eff} = N_f \ln[m^2 + (V_{12} + A_{12})(V_{21} + A_{21}) - im(\alpha + \beta) - \alpha\beta] \quad (38)$$

The overlaps of zero modes $V_{ij}(i, j = 1, 2)$ are of the order $V_{ij} \sim \rho^2 d^{-3}$ [1-3,16]. The leading correction to (38) in ρ/d stems from A_{ij} while that in $m\rho$ arises from α and β . Let us begin with the correction in ρ/d . For definiteness consider A_{12} . We are interested in the leading order correction and therefore can substitute K (see (25)) by S_0 . Since we are looking for the correction with respect to ρ/d we substitute Δ_i by δ_i (see (15)). Thus the quantity under consideration is

$$A'_{12} = \langle u^1 | S_0^{-1} \delta_2 S_0^{-1} \delta_1 S_0^{-1} | u^2 \rangle. \quad (39)$$

By chirality arguments $\delta_i(i = 1, 2)$ should not contain contributions proportional to $(1 + \gamma_5)/2$ (formally such contributions are found in $(S'_i - G_0)$, see (8)). Next we write

$$\begin{aligned} A'_{12} &\sim \int dz \left\{ \int dx u^{1+}(x) \partial_x \delta_2(x - z) \right\} \partial_z \left\{ \int dy \delta_1(z - y) \partial_y u^2(y - d) \right\} \equiv \\ &\equiv \int dz J_x(z) \partial_z J_y(z). \end{aligned} \quad (40)$$

Consider the contribution from the region $x \leq \rho, z \simeq d \pm \rho, y \simeq d \pm \rho$. In this region $u^{1+} \partial_x \sim \rho^{-3}$. To estimate $\delta_2(x - z)$ consider equation (8) for S'_2 with all arguments shifted by d since S'_2 corresponds to $\bar{I}$ placed at the distance d from the origin. It is easy to conclude that the leading contribution to $\delta_2(x - z)$ stems from the first term of the second line of (8) and $\delta_2(x - z) \sim G_0(x - z) \sim d^{-3}$. Therefore $J_x(z) \sim \rho^4 \rho^{-3} d^{-3} \sim \rho d^{-3}$. In $J_y(z)$ one has $\partial_y u^2(y - d) \sim \rho^{-3}$. The estimate of $\delta_1(z - y)$ deserves some attention. It also receives the main contribution from the first term of the second line of (18) but this time an additional small parameter arises:

$$\begin{aligned} \delta_1(z - y) &\sim G_0(z - y) [(\pi_x \pi_y)^{-1/2} (1 + \frac{\rho^2}{x^2 y^2} \tau_x \tau_y^+) - 1] \sim \\ &\sim G_0(z - y) \rho^3 d^{-3} \sim (z - y)^{-3} \rho^3 d^{-3}. \end{aligned} \quad (41)$$

Therefore

$$J_y(z) \sim \int dy (z-y)^{-3} d^{-3},$$

where at the final step the integration over y will yield the ρ^4 factor. Now ∂_z in (40) acting on $(z-y)^{-3}$ in (41) will result in $(z-y)^{-4} \sim \rho^{-4}$ since both z and y are $\sim (d \pm \rho)$. Substituting everything back into (40) we obtain

$$A'_{12} \sim \rho^5 d^{-6}. \quad (42)$$

Since V_{12} itself is $\sim \rho^2 d^{-3}$, the correction to it is of the order $0((\rho/d)^3)$.

Now we turn to $m\rho$ corrections. The leading one arises from α and β . Consider

$$\alpha' = \langle u^2 | S_0^{-1} \Delta_1 S_0^{-1} | u^2 \rangle. \quad (43)$$

Chirality arguments are important here. If we substitute both S_0^{-1} by G_0^{-1} and Δ_1 by δ_1 we get a zero. Therefore one should consider contributions arising from "unslashed" piece of any one of the three operators inside (43) and "slashed" pieces of the two others:

$$\begin{aligned} \alpha' \simeq & -im \langle u^2 | \delta_1 G_0^{-1} | u^2 \rangle - im \langle u^2 | G_0^{-1} \delta_1 | u^2 \rangle + \\ & + im \langle u^2 | G_0^{-1} (S_1'^2 - G_0^2) G_0^{-1} | u^2 \rangle \simeq im \langle u^2 | G_0^{-1} \delta_1^2 G_0^{-1} | u^2 \rangle. \end{aligned} \quad (44)$$

The last matrix element may be represented as

$$\begin{aligned} \alpha' \sim & m \int dz \left\{ \int dx u^{2+}(x-d) \partial_x \delta_1(x-z) \right\} \times \\ & \times \left\{ \int dy \delta_1(z-y) \partial_y u^2(y-d) \right\} \equiv m \int dz L_x(z) L_y(z). \end{aligned} \quad (45)$$

First consider the contribution from the "far" region $x, y, z \sim d \pm \rho$. Then $u^{2+}(x-d) \partial_x \sim \rho^{-3}$, $\delta_1(x-z) \sim d^{-3}$ (see (41)), $L_x(z) \sim \rho^4 \rho^{-3} d^{-3} \sim \rho d^{-3}$.

The same estimate holds for $L_y(z)$ and thus

$$\alpha' \sim m \rho^4 (\rho d^{-3})^2 \sim m \rho^6 d^{-6}. \quad (46)$$

One may similarly verify the estimate (46) for the "mixed" region $x, y \sim d \pm \rho, z \leq \rho$. Therefore the m^2 term in the effective action (38) acquires a correction of the order $0(\rho^6/d^6)$.

Thus, we conclude that the leading-order corrections to the effective action (38) are given by

$$\tilde{\Gamma}_{eff} = N_f \ln \{ m^2 [1 + 0(\rho^6/d^6)] + V_{12} V_{21} + (V_{12} + V_{21}) 0(\rho^5/d^6) \}, \quad (47)$$

and we remind that $V_{ij} \sim \rho^2/d^3$. We see that the corrections to $V_{12}V_{21}$ in the effective action are of order $\rho^3/d^3 \sim (1/3)^3$. A careful look into various normalization factors, powers of π , etc. enhances this result by an order of magnitude. Here however a caution is needed since we are not in a position to account for possible numerical factors resulting from the integration procedure.

Finally we wish to add one more remark on our results (38) and (47) for $\tilde{\Gamma}_{eff}$. The value of the overlap V_{ij} being proportional to ρ^2/d^3 depends also on the relative color orientation of I and $\bar{I}$. Maximal attraction is reached under the condition that the relative color orientation vector $u_\mu = tr(\Omega_{\bar{I}}\tau_\mu\Omega_I^+)$ is parallel to d_μ [14]. On the other hand the matrix element A_{ij} has a different dependence on the color orientation. Therefore the most favorable color orientations for $V_{ij}V_{ji}$ and $V_{ij}A_{ji}$ entering into $\tilde{\Gamma}_{eff}$ are different. The problem of the polarization of the $I\bar{I}$ molecule with higher modes included will be discussed elsewhere.

6 Summary

In this paper we have obtained exact solutions for the quark propagator and the effective action in the instanton–anti–instanton molecule. Up to now a consistent treatment of the non–zero–modes has been lacking. Possible applications of the results might be twofold. The first one concerns the scenarios of the chiral symmetry breaking and restoration involving $I\bar{I}$ molecules. Detailed properties of the fermion determinant are relevant for this subject – see e.g. [11–15]. Another possible application involves quark interactions induced by $I\bar{I}$ molecules and hadron correlators [14]. The corresponding studies were so far based on the quark propagator with non–zero–modes neglected. In such approximation the propagator exhibits the symmetry properties which are traceable in meson and baryon spectra. We have shown the relevance of non–zero–modes in the quark propagator. To which extent would higher modes affect the predicted properties of the spectrum? The results of the present paper allow to study this problem. The work is in progress now.

7 Acknowledgments

We wish to thank V.A.Rubakov for enlightening discussion which motivated our work. Fruitfull discussions with N.O.Agasyan and A.V.Smilga and especially Yu.A.Simonov are greatly acknowledged. The work was supported by the Russian Fund for Fundamental Research grants 97-02-16406, 96-02-19184, 96-15-96740 and RFFR-DFG 96-02-00088G.

References

- [1] T.Schäfer and E.V.Shuryak, Instantons in QCD, hep-ph/9610451.
- [2] D.Diakonov, Chiral symmetry breaking by instantons, in "Selected Topics in nonperturbative QCD", Varena 95, p.397.
- [3] M.A.Nowak, M.Rho, I.Zahed, Chiral nuclear dynamics, World Scientific, 1997.
- [4] M.C.Chu, J.M.Grandy, S.Huang, J.W.Negele, Phys. Rev. Lett. **70**, 225 (1993); Phys. Rev. **D49**, 6039 (1994).
- [5] B.O.Kerbikov, D.S.Kuzmenko, Yu.A.Simonov, JETP Lett. **65**, 128 (1997).
- [6] N.Andrei and D.J.Gross, Phys. Rev. **D18**, 468 (1978).
- [7] H.Levine and L.G.Yaffe, Phys. Rev. **D19**, 1225 (1979).
- [8] E.V.Shuryak and J.J.M.Verbaarschot, Nucl. Phys. **B410**, 37 (1993).
- [9] C.Callan, R.Dashen, and D.Gross, Phys. Lett. **63B**, 334 (1976).
- [10] C.Callan, R.Dashen, and D.Gross, Phys. Rev. **D17**, 2717, (1978).
- [11] M.-E.Ilgenfritz and E.V.Shuryak, Phys. Lett. **B319**, 511, (1989); *ibid.*, **B325**, 263 (1994).
- [12] V.V.Khoze and A.V.Yung, Z.Phys **C50**, 155, (1991).
- [13] Yu.A.Simonov, Yad. Fiz. **54**, 224, (1991).

- [14] T.Schäfer, E.V.Shuryak and J.J.M.Verbaarschot, Phys. Rev. **D51**, 1267 (1995).
- [15] T.Wettig, A.Schäfer, H.A.Weidenmüller, Phys. Lett. **B367**, 28 (1996).
- [16] C.Lee and W.A.Bardeen, Nucl. Phys. **B153**, 210 (1979).
- [17] M.Goldberger and K.M.Watson, Collision Theory, John Wiley, 1964.
- [18] B.O.Kerbikov, Theor. and Math. Physics, **65** 1225 (1985)
- [19] L.S.Brown, R.D.Carlitz, D.B.Creamer and C.Lee, Phys. Rev. **D17**, 1583 (1978).
- [20] G.'tHooft, Phys. Rev. **D14**, 3432, (1976).
- [21] D.I.Diakonov and V.Yu.Petrov, Nucl. Phys. **B272**, 457 (1986).
- [22] A.Vainshtein, V.Zakharov, V.Novikov and M.Shifman, Sov. Phys. Uspekhi **136**, 553 (1982).